

LiMPETS Rocky Intertidal Field Guide

Feather Boa Kelp (*Egregia menziesii*)



- Chocolate-brown to olive-green in color; up to 10 meters long.

- Long, flat stipe (stem) like a shoulder strap of a bag; fringed with small blades and floats for buoyancy.

- Young individuals have wide, textured axes (as above).

Sunburst Anemone (*Anthopleura sola*)



- Large, more than 2 inches (5 cm).

- Green to whitish in color; solitary.

- Has strong, clearly visible radiating lines on oral disk.

Giant Green Anemone (*Anthopleura xanthogrammica*)




- Large, more than 2 inches (5 cm).


- Olive-green to blue-green; solitary.

- Radiating lines on oral disk are absent or faintly visible.

Unidentified large, solitary anemone



- Large, more than 2 inches (5 cm); solitary. Closed enough so that oral disk is not visible.



- Body is greenish to white in color; often covered with shell debris.



- Do NOT count small (< 5 cm), closed anemones, as shown above.

Chitons (*Mopalia* spp./*Nuttallina* spp./*Tonicella* spp./others)

MOSSY CHITON



- Chitons are molluscs, oval in shape, with 8 overlapping shell plates.

CALIFORNIA SPINY CHITON



- Most are small, up to 2 inches (5 cm) wide.

LINED CHITON



- Often well camouflaged with surroundings.

Whelks (*Acanthinucella* spp./*Nucella* spp./others)

ANGULAR UNICORN SNAIL



- Whelks are predatory snails; shell aperture (opening) is typically oval.

EMARGINATE DOGWINKLE



- Shell is coiled or in a spiral; size and color vary.

CIRCLED ROCK SNAIL



- Both ends of shell are pointed.

Turban Snails (*Chlorostoma brunnea/funebralis*)



BLACK TURBAN SNAIL

- Up to 1 inch (2.5 cm) long. Color deep purple, black, or brown.



BLACK TURBAN SNAIL

- Always check to make sure it is a snail and not a hermit crab.



BROWN TURBAN SNAIL

- Shell is smooth, a rounded cone shape (no point at tip of shell).

Hermit Crabs (*Pagurus* spp.)



HAIRY HERMIT CRAB

- All hermit crabs use snail shells as portable homes. The one above has white bands on its walking legs.



GRAINYHAND HERMIT CRAB


- Some (as above) have walking legs with light blue/white flecks; solid red antennae.



BLUEBAND HERMIT CRAB

- Some (as above) have walking legs with blue bands at the tips; solid red antennae.

Purple Sea Urchin (*Strongylocentrotus purpuratus*)



- Up to 4 inches (10 cm) in length; reddish to purple in color.



- Juveniles are pale green. Individual above approximately the size of a nickel.




- Spherical body covered with spines. Often rocks and shells attached.

Green Pin-cushion Alga (*Cladophora columbiana*)



- Bright green and spongy.



- Consists of branched filaments that form densely matted tufts.



- Resembles clumps of moss.

Dead Man's Fingers (*Codium fragile*)



- Can be fairly large, up to 16 inches (40 cm) in length.



- Dark green to blackish-green in color.



- The “fingers” are forked, spongy and are about as thick as a pencil or pen.

Sea Lettuces (*Ulva* spp.)



- Oval shaped blades, up to 16 inches (40 cm); bright green or yellow-green.




- Usually grow as sheets, but one species exists in a cylindrical form (as above).



- Thin, almost transparent sheets, only 2 cell layers thick; often look like wilted lettuce.

Surfgrasses (*Phyllospadix scouleri/torreyi*)



- Up to 0.5 cm wide and 6.5 feet (2 m) in length; flowers are small, inconspicuous.
- Leaves are bright green, narrow, long and wiry.
- Photo above shows a close-up of the female flower stalk with seeds.

Flattened Rockweeds (*Fucus gardneri/Hesperophycus californicus*)



- Can be olive-green to tan in color; up to 10 inches (25 cm) tall.
- Reproductive tips can often be swollen.
- Flattened body, wide blades with distinct midrib, dichotomous branching.

Slender Rockweeds (*Pelvetiopsis limitata/Silvetia compressa*)



- Can be olive-green to tan in color; 2-35 inches (5-90 cm) tall.
- Can be darker, shriveled and tough when dried out.
- Flattened body, thin blades with NO midrib, dichotomous branching.


Tar Spot Algae (*Mastocarpus* spp./*Ralfsia* spp./others)

RALFSIA SP.



- Black crust on rock, looks like tar.

MASTOCARPUS SP. & RALFSIA SP.



- Some feel rough. Others feel thicker, more spongy.

MASTOCARPUS SP.



- Can grow in small or large patches (as above).


Encrusting Coralline Algae (many species)



- Light or bright pink crust on rock.



- Crust smooth or covered in bumps.



- Can grow in small or large patches.

Upright Coralline Algae (*Bossiella* spp./*Calliarthron* spp./*Corallina* spp.)

CORALLINA SP.



- Light whitish-pink to bright pink in color.

CALLIARTHRON SP.



- Calcium carbonate in cell walls can make them feel relatively stiff.

CORALLINA SP.



- Many species are branched and have tiny, jointed segments.

Scouring Pad Alga (*Endocladia muricata*)



- Short, bushy clumps; 1-3 inches (3-8 cm) tall.



- Dark reddish-brown in color.



- Branches covered with short spines (feels rough, not slimy or smooth).

Stunted Turkish Towel (*Mastocarpus* spp./*Mazzaella affinis*)

MASTOCARPUS JARDINII



- Species above is red, brown to blackish; up to 4 inches (10 cm) tall. Blades are narrow and have bumps.

MASTOCARPUS PAPILLIATUS



- Species above is light red to purplish-black. Blades are wider, split at the tips, often have bumps.

MAZZAELLA AFFINIS



- Species above is olive to reddish-brown; up to 6 inches tall (15 cm). Blades smooth.

Lawn Alga (*Chondracanthus canaliculatus*)



- Low and bushy with flat but sharp, pointed tips; up to 6 inches (15 cm).



- Yellow-green in warm waters to olive-purple in colder waters.



- Forms mat-like clumps of entangled branches (feels smooth, not rough). Can be mixed in with other algae.

Nori (*Porphyra* spp.)



- Color varies from brown, yellow-green to purple. Can resemble sea lettuce but nori is NOT bright green.

Blades are often ruffled at the edges; only 1-2 cell layers thick. Size varies.

- Can resemble crumpled cellophane when dry.

Sea Sacs (*Halosaccion glandiforme*)




- Up to 6 inches (15 cm) tall.

- Yellowish-brown, hollow sacs usually filled with seawater.

- As the alga gets older, the tips of the sacs can erode and leave the alga flat or filled with sand.

Iridescent Algae (*Mazzaella flaccida/splendens*)

MAZZAELLA SPLENDENS



MAZZAELLA SPLENDENS



MAZZAELLA FLACCIDA



- Large, oval or heart shaped blades; up to 12 inches (30 cm) tall.

- Can appear iridescent; dark purple, brown or green in color.

- Can also appear as above; yellowish-green blade with purple or brown near base of blade,

Aggregating Anemones (*Anthopleura elegantissima*)



- Small, less than 2 inches (5 cm); often with pink-tipped tentacles.



- Greenish body.



- Can form dense aggregations: often covered in sand and shells.

Honeycomb Tube Worm (*Phragmatopoma californica*)



- Also called sandcastle worms; live in tubes of cemented sand grains, each with a flared rim.



- Often in large masses up to 6.5 feet (2 m) in length.



- Tubes regularly placed in a honeycomb arrangement.

Limpets (*Lottia* spp.)

L. SCABRA, ROUGH LIMPET



- Snail-like mollusc with one shell. Most are < 1 inch (2.5 cm).

L. DELTA, SHIELD LIMPET



- Shell is cone shaped or flat; smooth or ribbed in texture.

L. GIGANTEA, OWL LIMPET



- Owl limpet above can grow to 4 inches (10 cm).

Sea Mussel (*Mytilus californianus*)



- Shell up to 8 inches (20 cm); bluish-black in color; radial ribbing.



- Bivalve mollusc with two shells; use byssal threads to attach to rock.



- Can form extensive beds that create habitat for many species.

Leaf Barnacle (*Pollicipes polymerus*)



- Also called gooseneck barnacles; up to 3 inches (8 cm) in length.



- Strong, dark brown, rubbery stalk; topped with 5 or more white plates.



- Usually found in tight clusters; often mixed with sea mussels.

Common Acorn Barnacles (*Balanus glandula*/*Chthamalus dalli*/*fissus*)



- Small in size; shell up to $\frac{3}{4}$ inch (2 cm) wide.



- Shell white or brownish.



- Ribbed or smooth outer plates.

Pink Acorn Barnacle (*Tetraclita rubescens*)



- Large barnacle, up to 2 inches (5 cm) wide.



- Shell is reddish-pink, appearing thatched.




- **Do NOT count the above barnacle; similar in size but whitish-brown in color.**


Bare Rock



- Bare, rocky substrates larger than sand or gravel.



- Contains no obvious living organisms (as in circle above).



- Even small patches of bare rock within square(s) should be counted.

Loose Sand



- Granular (fine sand to gravel) substrate.



- Sand must be loose, unattached to anemones or other organisms.



- Even small patches of sand within square(s) should be counted.

Tidepool and Monitoring Etiquette



- Avoid stepping on invertebrates and algae whenever possible.
- Return animals where you found them.
- Replace rocks where you found them.
- Do not take anything except pictures.
- Do not leave any trash at the site.
- Avoid wading in tidepools.
- Always keep an eye on the water and don't turn your back on the ocean, even for a moment.
- Be aware of your surroundings, including the water, slippery rocks or algae, and tidepool creatures.

Acknowledgements

Funder: Friends of Fitzgerald Marine Reserve, <http://fitzgeraldreserve.org>

Design, Development and Editing:

Stuart Chan, Ison Design, www.isondesign.com

Abby Nickels & Amy Dean: Farallones Marine Sanctuary Association & LiMPETS, <http://limpetsmonitoring.org>

LiMPETS Science Advisory Panel: Steven Lee (UCLA), Steve Lonhart (SIMoN/MBNMS), John Pearse (UCSC/PGMNH)

Photo Credits: *We gained permission from the following photographers, scientists, and educators to use their photos in this field guide. Many thanks for their generosity. Photos may not be used, altered or duplicated without their permission.*

Folini, Franco: Photographer, www.flickr.com/photos/78425154@N00/5789895873

Janiak, Courtney: Photographer, www.flickr.com/photos/seaweedlady

King, Chad: SIMoN/MBNMS, www.sanctuariesimon.org/photos/index.php

Kirkhart, Jerry: Photographer, www.flickr.com/people/jkirkhart35

Lee, Steven: Research Coordinator, Aquatic Biologist, UCLA

LiMPETS staff: Altstatt, J., Dean, A.F., Emanuelson, L., Fisher, A., Nickels, A., Wasser, A. and A. Young

Lonhart, Steve: SIMoN/MBNMS, www.sanctuariesimon.org/photos/index.php

McDonald, Gary: Long Marine Lab, UCSC, <http://calphotos.berkeley.edu>

Mosquin, Daniel: UBC Botanical Garden, www.botanicalgarden.ubc.ca

Oregon State Parks: <http://www.flickr.com/photos/orstateparks/5229773010/in/photostream>

Pacific Coast Rocky Intertidal Program: www.eeb.ucsc.edu/pacificrockyintertidal/index.html

Pearse, John: UCSC/Pacific Grove Museum of Natural History (LiMPETS)

Pederson, Josh: SIMoN/MBNMS, www.sanctuariesimon.org/photos/index.php

Ueda, Ken-Ichi: iNaturalist.org, www.flickr.com/people/ken-ichi

Watanabe, James: Hopkins Marine Station, Stanford Univ., <http://seanet.stanford.edu>

Wood, Wendell: algaeBASE, www.algaebase.org

Young, Alison: California Academy of Sciences, www.calacademy.org

